



# General Considerations in the Formulation and Evaluation of Herbal Hair Oils

1. Harshal Dattatraya Amale 2. Vijay Kumar S. Wakale 3. Gandharva Anil Marne

1. Student, Samarth Institute of Pharmacy, Belhe, Pune, Maharashtra, India.
2. Professor, Department of Pharmaceutical Chemistry, Samarth Institute of Pharmacy, Belhe, Pune, Maharashtra, India
3. Student, Samarth Institute of Pharmacy, Belhe, Pune, Maharashtra, India.

## How to Cite this Article:

Amale, H. D. & Marne, G. A. (2026). General Considerations in the Formulation and Evaluation of Herbal Hair Oils. International Journal of Creative and Open Research in Engineering and Management, <i>02</i>(05).  
<https://doi.org/10.55041/ijcope.v2i5.681>

## License:

This article is published under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

© The Author(s). Published by International Journal of Creative and Open Research in Engineering and Management.



<https://doi.org/10.55041/ijcope.v2i5.681>

## Abstract

A particular connection between lipid chemistry and plant medicines is represented by the pharmacological production of herbal hair oils. With an emphasis on carrier selection, bioactive extraction efficiency, and analytical validation, this review offers a thorough summary of the general factors necessary for formulating high-efficacy herbal hair oils. The scope includes a critical assessment of vegetable lipids such as Sesamum indicum and Cocos nucifera as delivery vehicles, highlighting their potential to interact with the 18-methyl eicosanoic acid (18-MEA) layer and enter the hair cuticle. The ability of important production techniques, particularly the conventional heat-infusion method to simultaneously absorb fat-soluble and water-soluble secondary metabolites is examined. Additionally, the manuscript describes the molecular mechanisms of action, such as the stability of beta-catenin within the pilosebaceous unit and the inhibition of 5-alpha reductase. Physicochemical benchmarks including acid value, saponification value, and refractive index are used to address analytical standardization in addition to stability issues pertaining to lipid oxidation kinetics. This review concludes important gaps in regulatory harmonization and the integration of cutting-edge drug delivery systems, like nanoemulsions, by reviewing recent literature. This provides a path for the manufacturing of herbal hair care preparations that are safe,

effective, and repeatable for the global market.

**Keywords:** Taila Paka Vidhi, 5-alpha Reductase, 18-MEA Layer, Herbal Hair Oil, Lipid Bioavailability, Physicochemical Standardization, Lipid Oxidation, Follicular Targeting, Anagen Phase, Pharmaceuticals.

## Introduction

In today's pharmaceutical environment, hair is recognized not only as an attractive characteristic but also as a complex biological indicator of systemic health and a primary barrier against environmental stresses. The human hair follicle (HF) is a unique, tightly regulated mini-organ that goes through cycles of growth (anagen), regression (catagen), and rest (telogen). Different forms of alopecia, which affect about 2% of people globally, are brought on by disturbances to this cycle that are amplified by factors like oxidative stress, androgenic activity, and environmental pollutants.



The development of herbal hair oils is being driven by a global paradigm shift toward Herbal Drug Technology, as consumers look for natural alternatives to synthetic agents like finasteride and minoxidil. Herbal hair oils are multifunctional delivery systems that provide essential nutrients including flavonoids, polyphenols, and vitamins that support sebaceous gland activity while also providing mechanical protection for the hair shaft.

The way a hair oil interacts with the 18-methyl eicosanoic acid (18-MEA) layer, the outermost hydrophobic barrier on the hair shaft, determines its efficacy from a physicochemical perspective. Effective herbal oils must penetrate the cuticle in order to reduce "hygral fatigue" and prevent protein loss; this capacity varies significantly depending on the lipid chain length of the carrier oil. These preparations have their roots in the Ayurvedic branch of Bhaishajya Kalpana, namely Sneha Kalpana, where long-boiling medicinal oils transfer bioactive components into a stable lipid matrix. It is currently important from a pharmacological standpoint to standardize these traditional extraction methods to ensure therapeutic reproducibility. Temperature and duration must be carefully regulated to prevent the degradation of heat-sensitive alkaloids and essential oils.

Furthermore, current research is increasingly concentrating on the molecular signaling pathways like the beta-catenin and IGF-1 pathways through which herbal actives stimulate follicular regeneration. The purpose of this paper is to provide an analytical overview of the production procedures, quality assurance requirements, and formulation components needed to transform traditional herbal remedies into standardized pharmaceutical products. By compiling data from over 40 references, this publication offers researchers and industrial formulators a crucial foundation to optimize the creation of herbal hair oils.

### Plant profile

Common Name	Botanical Name	Family	Part Used	Important Phytoconstituents
<b>Amla</b>	<i>Phyllanthus emblica</i>	Phyllanthaceae	Fruit	Vitamin C, tannins, gallic acid, ellagic acid
<b>Bhringraj</b>	<i>Eclipta alba</i>	Asteraceae	Whole plant	Wedelolactone, ecliptine, flavonoids
<b>Neem</b>	<i>Azadirachta indica</i>	Meliaceae	Leaves	Nimbin, azadirachtin, quercetin
<b>Hibiscus</b>	<i>Hibiscus rosa-sinensis</i>	Malvaceae	Flowers	Flavonoids, anthocyanins, mucilage,

### Materials

Component Type	Specific Ingredient(s)	Role in Formulation
Primary Base	Coconut Oil ( <i>Cocos nucifera</i> )	Main oil base and carrier
Secondary Base	Castor Oil ( <i>Ricinus communis</i> )	Supports the base and improves texture



Active Ingredient	Bhringraj ( <i>Eclipta alba</i> )	Promotes hair growth
Nutrient Active	Amla ( <i>Phyllanthus emblica</i> )	Nourishes and protects hair
Scalp Purifier	Neem ( <i>Azadirachta indica</i> )	Cleanses scalp and controls infection
Conditioning Agent	Fenugreek ( <i>Trigonella foenum-graecum</i> )	Softens hair and improves shaft strength
Aromatic Stimulant	Rosemary Oil	Improves scalp stimulation
Antioxidant	Vitamin E ( $\alpha$ -Tocopherol)	Protects formulation from oxidation

### Equipments

1. **Brookfield Viscometer** – Used to measure viscosity and study the flow behavior of the formulation.
2. **Abbe Refractometer** – Measures refractive index to check purity and consistency of the oil base.
3. **Digital pH Meter** – Determines pH to ensure compatibility with the scalp's natural acidic pH.
4. **Specific Gravity Bottle** – Used to determine the density and weight per mL of the formulation.
5. **HPTLC System** – Helps identify and quantify important phytochemical constituents in the formulation.

Ingredient	Quantity (w/w)	Percentage (%)
<b>Amla Extract (Heat Infusion)</b>	1.0 g	10.0
<b>Bhringraj Powder (Coarse)</b>	1.0 g	10.0
<b>Neem Leaves (Fresh/Dried)</b>	0.5 g	5.0
<b>Hibiscus Flowers (Dried)</b>	0.5 g	5.0

<b>Castor Oil</b>	1.5 g	15.0
<b>Coconut Oil</b>	q.s. 10.0 g	55.0



## Formulation Table

### Method of preparation

1. All instruments required for the preparation were checked, calibrated, and the apparatus was properly cleaned before starting the experiment.
2. The selected fresh or dried herbal materials were weighed and crushed to obtain a coarse paste known as Kalka.
3. A herbal decoction (Kwatha) was prepared by boiling the plant materials in water until the liquid was reduced to approximately one-fourth of its original volume.
4. The base oil was subjected to Taila Murchhana by heating it with suitable herbal ingredients to reduce moisture content and improve the quality of the oil.
5. The prepared oil, herbal paste, and decoction were mixed together in the desired proportion in a suitable vessel.
6. The mixture was heated at a controlled temperature with continuous stirring to allow proper incorporation of herbal constituents into the oil.
7. The progress of the preparation was observed regularly for characteristic signs of Sneha Siddhi, including reduction of frothing and disappearance of crackling sound.
8. A small quantity of the herbal paste was tested between the fingers to check whether it could be rolled smoothly into a wick-like form without sticking.
9. The prepared oil was filtered in hot condition using a clean muslin cloth to separate unwanted residues.
10. After cooling the formulation below 40°C, essential oils and Vitamin E were incorporated and mixed thoroughly.
11. The final herbal hair oil was filled into amber-coloured glass containers and stored away from direct light exposure.

### Evaluation parameters

Parameter	Acceptable Range	Significance
Acid Value	≤ 4.5	Acid value shows the amount of free fatty acids present in the A higher value may indicate hydrolysis and early rancidity.
Saponification Value	19.15 – 224.2	Saponification value helps estimate the average molecular weight of fatty acids present in formulation.



<b>Specific Gravity</b>	0.83 – 1.09	Specific gravity is measured to confirm the density, purity, and batch consistency of the formulation.
<b>Refractive Index</b>	1.29 – 1.50	Refractive index is used as a quality control parameter to evaluate the degree of unsaturation in oils.
<b>Viscosity (cps)</b>	28 – 35	Viscosity determines the flow property, spreadability, and overall application feel of the formulation.
<b>pH</b>	5.5 – 7.3	Maintaining this pH range helps keep the formulation compatible with the natural acidic pH of the scalp.

## Result

According to the review, coconut oil's low molecular weight allows it to deeply penetrate the hair cortex, making it one of the best carrier oils. The heat-infusion approach (Taila Paka) is thought to be more successful during formulation preparation since it enables better extraction of both oil-soluble and water-soluble phytoconstituents. The formulation must be properly standardized, and quality and stability are frequently verified using metrics like specific gravity (0.83–1.09) and acid value ( $\leq 4.5$ ). By altering Wnt/ $\beta$ -catenin signaling pathways and blocking 5-alpha reductase (5AR), which is linked to follicular damage and hair loss, the active herbal components may promote hair growth.

## Reference

1. Marina AM, Man YC, Nazimah SA, Amin I. Chemical properties of virgin coconut oil. *J Am Oil Chem Soc.* 2009;86:301–307.
2. Shukla P, Joshi D, Kumar Y, Chaturvedi M. Herbal hair oil for enhancement of hair growth: formulation and evaluation. *Int J Pharm Res Dev.* 2026;8:12–14.
3. Sarkar R, Podder I, Gokhale N, Garg VK. Use of vegetable oils in dermatology: an overview. *Int J Dermatol.* 2017;56:1080–1086.
4. Dixit VK, Adhirajan N, Gowri C. Development and evaluation of herbal formulations for hair growth. *Indian Drugs.* 2001;38:559–563.
5. Adhirajan N, Ravikumar T, Babu M. In vivo and in vitro evaluation of hair growth potential of *Hibiscus rosa-sinensis* Linn. *J Ethnopharmacol.* 2003;88:235–239.
6. Patni P, Varghese D, Jain DK. Formulation and evaluation of herbal hair oil for alopecia management. *Planta Indica.* 2006;2:27–30.
7. Banerjee PS, Sharma M, Nema RK. Preparation, evaluation and hair growth stimulating activity of herbal hair oil. *J Chem Pharm Res.* 2009;1:261–267.
8. Jaiswal S, Tiwari S, Chaturvedi M, Gupta R. Formulation, development and evaluation of herbal hair oil. *Diary Med Conveyance Ther.* 2017;7:178–179.
9. Joshi AA, Dyawarkonda PM. Formulation and evaluation of polyherbal hair oil. *Int J Green Pharm.* 2017;11:S135–S138.
10. Oyi AR, Onaolapo JA, Obi RC. Formulation and antimicrobial studies of coconut (*Cocos nucifera* Linne) oil. *Res J Appl Sci Eng Technol.* 2010;2:133–137.
11. Keis K, Persaud D, Kamath YK. Investigation of penetration of coconut oil into human hair by SIMS. *J Cosmet Sci.* 2005;56:283–295.
12. Valamuri H, Lavanya C, Naga H. Herbal goodness of hair: a comprehensive review. *Int J Pharm Sci.* 2024;2:1851–1865.
13. Khanpara P, Sorathiya A, Faldu S. A review article on the formulation and evaluation of herbal hair oil. *IJPRA J.* 2024;9:1120–1128.



14. Singh BK, Roy S. Preparation and evaluation of Ayurvedic medicated oil (Sneha Kalpana): a critical review. *RJPP*. 2016;8:45–52.
15. Bagad YM, Barhate SD, Shaikh SK. Design, formulation and evaluation of herbal hair oil for enhance hair growth activity. *RJTCS*. 2025;16:1–9.
16. Cho YH, Lee SY, et al. Effect of pumpkin seed oil on hair growth in men with androgenetic alopecia. *Evid Based Complement Alternat Med*. 2014;2014:549721.
17. Datta K, Singh AT, et al. *Eclipta alba* extract with potential for hair growth promoting activity. *J Ethnopharmacol*. 2009;124:450–456.
18. Roy RK, Thakur M, Dixit VK. Hair growth-promoting activity of *Eclipta alba* in male albino rats. *Arch Dermatol Res*. 2008;300:357–364.
19. Mahawar V, Mahawar S. Formulation development and evaluation of herbal hair oil. *ResearchGate*. 2024.
20. Rahane RD, et al. To design, formulate and evaluate herbal hair growth oil. *J Adv Zool*. 2024;45(2):1791–1805.
21. Sharma R, Amin H, Prajapati PK. Pharmaceutical standardization of Shadbindu Taila. *PMC*. 2013;PMC3764865.
22. Mourya ND, Kare V. A comprehensive review of formulation and evaluation of herbal hair oil. *IJPSJ*. 2024;9:1851–1865.
23. Gandhi SM, Swarn S, More UA. Liposomal systems for targeted drug delivery to hair follicles. *J Drug Deliv Sci Technol*. 2025;107624.
24. Khairnar TD, Chavan GS, Agrawal YO. Recent trends in nanocarrier formulations in androgenetic alopecia management. *J Drug Deliv Sci Technol*. 2024;98:105890.
25. Kalompatsios D, et al. Evaluation of oxidative stability and antioxidant capacity of infused olive oil. *Lipidology*. 2026;3:14.
26. Singh AP, et al. Antioxidants help favorably regulate the kinetics of lipid peroxidation in hempseed oil. *PMC*. 2020;PMC7324387.
27. Panizzon GP, Vercellino AC. On hair care physicochemistry: from structure to conditioning agents. *PMC*. 2023;PMC9921463.
28. Cho S, Kim S, Park H, et al. Phytochemicals targeting hair growth-related molecular pathways. *PMC*. 2025;PMC12603380.
29. Rele AS, Mohile RB. Effect of mineral oil, sunflower oil, and coconut oil on prevention of hair damage. *J Cosmet Sci*. 2003;54:175–192.
30. *Ayurvedic Pharmacopoeia of India*. Vol. II. New Delhi: Government of India; 2014.
31. Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. Pune: Nirali Prakashan; 2008.
32. *Indian Pharmacopoeia*. Vol. I. Ghaziabad: Indian Pharmacopoeia Commission; 2010.
33. Suresh Kumar P, et al. In vitro evaluation of anti-dandruff activity of polyherbal hair oil. *J Pharm Res*. 2010;3:2956–2958.
34. Tiwari G, Tiwari R. Assessment of nutraceutical potential of herbs for promoting hair growth. *RJPP*. 2021;13:150–162.
35. Azeem A, et al. Nanoemulsion components screening and selection: a technical note. 2018.
36. Cardoso S, Barradas TN. Nanoemulsions loaded with minoxidil and clove oil for drug follicular targeting. *J Drug Deliv Sci Technol*. 2020.
37. Deshpande DJ. *Handbook of Herbal Remedies*. 2008.
38. Saleem K, Siddiqui B. Minoxidil nano-hydrogel based on chitosan and hyaluronic acid. *J Drug Deliv Sci Technol*. 2020.



39. Choe E, Min DB. Mechanisms and factors for edible oil oxidation. *J Food Sci.* 2006;71:R172–R186.
40. Sabarwal N, et al. Development and evaluation of polyherbal formulations for hair growth activity. *PHCOG J.* 2009;1(2):165–170.